PHAP Antibody

Catalog No: #24199

Description



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name PHAP Antibody Host Species Rabbit Clonality Polyclonal Purification Affinity chromatography purified via peptide column E WB IHC Applications Species Reactivity Hu Ms Rt Peptide Immunogen Type Immunogen Description PHAP antibody was raised with a synthetic peptide corresponding to amino acids at carboxy terminus of human PHAP I. Target Name PHAP Accession No. P39687 Formulation Supplied in PBS containing 0.02% sodium azide. Storage Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Application Details

Predicted MW: 35 kd

Images



Western blot analysis of PHAP expression in human Raji cell Ilysate with PHAP antibody at 1 ug/mL.



Immunohistochemistry of PHAP in human small intestine tissue with PHAP antibody at 10 ug/mL.

Background

Apoptosis is related to many diseases and development. Caspase-9 plays a central role in cell death induced by a variety of apoptosis activators. Cytochrome c, after released from mitochondria, binds to Apaf-1, which forms an apoptosome that in turn binds to and activate procaspase-9. Activated caspase-9 cleaves and activates the effector caspases (caspase-3, -6 and -7), which are responsible for the proteolytic cleavage of many key proteins in apoptosis. The tumor suppressor putative HLA-DR-associated proteins (PHAPs) were recently identified as important regulators of mitochondrion apoptosis. PHAP appears to facilitate apoptosome-medicated caspase-9 activation and to stimulate the mitochondrial apoptotic pathway. PHAP was also shown to oppose both Ras- and Myc-medicated cell transformation.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.