TIRAP Antibody

Catalog No: #24201




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

_					
	esc	٦rı	nt	in	n
		7	О.	II J	

Product Name	TIRAP Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Affinity chromatography purified via peptide column	
Applications	E WB IHC	
Species Reactivity	Hu Ms Rt	
Immunogen Type	Peptide	
Immunogen Description	Raised against a peptide corresponding to 15 amino acids near the C-terminus of mouse TIRAP.	
Target Name	TIRAP	
Accession No.	AAL05628	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	

Images



Western blot analysis of TIRAP in human heart cell lysates with TIRAP antibody at (A) 0.5, (B) 1, and (C) 2 ug/mL.



Immunohistochemical staining of human heart cells using TIRAP antibody at 2 ug/mL.

Background

Toll-like receptors (TLRs) are signaling molecules that recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. These proteins act through adaptor molecules such as TIRAP and MyD88 to activate various kinases and

transcription factors. In TIRAP-deficient mice, TLR signaling in response to TLR2 ligands (using either TLR1 and TLR6 as co-receptors) is totally abolished, suggesting that MyD88 and TIRAP work together and are both required for TLR2 signaling. Furthermore, these mice are also resistant to the toxic effects of LPS and show defects in NF-?B and MAP kinase activation, suggesting that TIRAP is also involed in TLR4 signaling.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.