GITRL Antibody

Catalog No: #24341


Description




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name GITRL Antibody Host Species Rabbit Clonality Polyclonal Purification Affinity chromatography purified via peptide column E WB ICC Applications Species Reactivity Hu Ms Recombinant protein Immunogen Type Immunogen Description Raised against purified recombinant human GITR ligand. Target Name GITRL Other Names AITRL Accession No. Q9UNG2 Formulation Supplied in PBS containing 0.02% sodium azide. Storage Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of (A) 5 ng, (B) 25 ng, and (C) 50 ng of purified recombinant GITRL with ITRL antibody at 1 ug/mL.



Immunocytochemistry of GITRL in THP-1 cells with GITRL antibody at 10 ug/mL.

Background

The tumor necrosis factor (TNF) and TNF receptor (TNFR) gene superfamilies regulate numerous biological functions including cell proliferation, differentiation, and survival through regulating the activation of the transcription factor NF-kB and various mitogen-activated protein kinases. The glucocorticoid-induced tumor necrosis factor receptor (GITR) is an emerging member of this family that is expressed on CD4+ CD25+ regulatory T cells and appears to have crucial immune regulation functions. Its ligand GITRL is expressed in endothelial and antigen-presenting cells and can activate NF-kB, induce both pro- and anti-apoptotic effects, inhibit the suppressive activity of regulatory T cells, and co-stimulate responder T cells through GITR. Dominant negative forms of NIK and TRAF2 expressed in transfected 293 cells substantially inhibited NF-kB activation, suggesting that the GITRL-GITR pathway involves both NIK and TRAF2.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.