AES Antibody

Catalog No: #24349




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			
	escri	nt	ION.
\boldsymbol{L}	COUL	IJΨ	ULL

Product Name	AES Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Affinity chromatography purified via peptide column	
Applications	E WB ICC	
Species Reactivity	Hu Ms Rt	
Immunogen Type	Peptide	
Immunogen Description	Raised against a 16 amino acid peptide from near the carboxy terminus of human AES.	
Target Name	AES	
Other Names	Amino-terminal enhancer of split, GRG, ESP1, TLE5	
Accession No.	NP_945320	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	

Images



Western blot analysis of AES in 293 cell lysate with AES antibody at (A) 1, (B) 2 and (C) 4 μ mL.



Immunocytochemistry of AES in HeLa cells with AES antibody at 10 ug/mL.

Background

Adhesion to extracellular matrix regulates cell survival through both integrin engagement and appropriate cell spreading. Anoikis is the molecular mechanism of apop-tosis induced by integrin detachment. Amino-terminal enhancer of split (AES) is a member of the Groucho/ transducin-like enhancer of split (TLE) family of transcriptional regulators, a group of transcriptional co-repressors that play important roles in neurogenesis, segmentation, and sex determination. AES forms a complex with Bit1 (Bcl-2 inhibitor of transcription 1), a mitochondrial protein that is released into the cytoplasm upon onset of apoptosis. It has been suggested that this complex turns off a survival-promoting gene transcription program controlled by the TLE protein family. Interestingly, apoptosis of cells transfected with AES and Bit1 could be inhibited if the cells were allowed to attach to fibronectin through the alpha5beta1 integrin suggesting that the Bit1-AES pathway contributing to anoikis is regulated by integrins, and in particular, the alpha5beta1 integrin.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.