GAPDH Antibody

Catalog No: #24407




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			
	escr	מוי	tion
\boldsymbol{L}	COUL	ıv	เเบเ

Product Name	GAPDH Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Affinity chromatography purified via peptide column	
Applications	E WB IHC	
Species Reactivity	Hu Ms Rt	
Immunogen Type	Peptide	
Immunogen Description	Raised against a 16 amino acid peptide from near the amino-terminus of human GAPDH.	
Target Name	GAPDH	
Other Names	Glyceraldehyde-3-phosphate dehydrogenase, G3PDH, GAPD	
Accession No.	P04406	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	

Images



Western blot analysis of GAPDH in HeLa cell lysate with GAPDH antibody at (A) 0.125, (B) 0.25 and (C) 0.5 ug/mL.



Immunohistochemistry of GAPDH in human liver tissue with GAPDH antibody at 10 ug/mL.

Background

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD), an important energy-yielding step in carbohydrate metabolism. Recent evidence suggests that it also is involved in a number of cellular processes such as membrane fusion, phosphotransferase activity, DNA replication and repair, and nuclear RNA export. GAPDH has also been implicated in playing a role in different pathologies such as cancer progression, apoptosis, and neuronal diseases such as Alzheimerß s and Huntingtonß s disease. GAPDH is constitutively expressed at high levels in almost all tissues and cell lines making it ideal for use as a loading control marker in immunoblots.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.