erbB-2 Antibody

Catalog No: #24888




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_				
	esc	rır	۱t17	nn.
		7 1 1 4	лι	7/1

Product Name	erbB-2 Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Affinity chromatography purified via peptide column	
Applications	E WB IHC	
Species Reactivity	Hu Ms Rt	
Immunogen Type	Peptide	
Immunogen Description	Raised against a 19 amino acid peptide near the carboxy terminus of human erbB-2.	
Target Name	erbB-2	
Other Names	erbB-2, v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, NEU, NGL, HER2, HER-2, TKR1, CD340	
Accession No.	NP_004439	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	

Images



Western blot analysis of erbB-2 in rat liver tissue lysate with erbB-2 antibody at (A) 1 and (B) 2 ug/mL.



Immunohistochemistry of erbB-2 in human liver tissue with erbB-2 antibody at 2.5 $\mbox{ug/mL}.$

Background

The ErbB family consists of four closely related tyrosine kinase receptors that act as potent mediators of normal cell growth and development. Aberrant expression or function of one or more of these receptors can play a major role in the development and evolution of cancer. ErbB-2, also known as HER2, has been implicated in the evolution of both breast and gastric cancers, and is evident in other cancer types such as ovarian and salivary gland tumors. ErbB-2 possesses an active tyrosine kinase domain, but no direct ligand has been identified yet. ErbB-2 is the preferred binding partner to the other members of the ErbB family and is thought to act primarily through the Ras-MAPK, Pl3k-PKB/Akt, and PLC-PKC signaling pathways. Numerous anti-cancer strategies have been employed against erbB-2, such as antibody-based therapies to prevent ligand binding or receptor activation through dimerization, antibody-dependent cell mediated cytotoxicity, in addition to direct kinase inhibition to prevent molecular activation/downstream signaling.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.