LIF Antibody

Catalog No: #25283



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description	-111
Product Name	LIF Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	E WB
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide
Immunogen Description	Raised against a 16 amino acid peptide near the center of human LIF.
Target Name	LIF
Other Names	Leukemia inhibitory factor, CDF, DIA, HILDA, MLPLI
Accession No.	NP_002300

Supplied in PBS containing 0.02% sodium azide.

Images

Formulation

Storage



Western blot analysis of LIF in 3T3 cell lysate with LIF antibody at (A) 1 and (B) 2 ug/mL.

Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated

freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Background

LIF is a pleiotropic cytokine with roles in several different systems. It is involved in the induction of hematopoietic differentiation in normal and myeloid leukemia cells, induction of neuronal cell differentiation, regulator of mesenchymal to epithelial conversion during kidney development, and may also have a role in immune tolerance at the maternal-fetal interface. LIF was initially recognized by its ability to induce terminal differentiation of myeloid leukemic cells. It is a member of the IL-6 cytokine superfamily and can be highly glycosylated. LIF signaling is transduced through the LIF-R/gp130 receptor complex, leading to the phosphorylation and activation of the JAK/STAT pathway. Recent evidence shows that LIF inhibits cardiomyogenesis in embryonic stem cells via STAT3 activation.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.