

CBFB Antibody

Catalog No: #32759

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	CBFB Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were purified by affinity purification using immunogen.
Applications	WB IHC IF
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total CBFB protein.
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant protein of human CBFB.
Target Name	CBFB
Other Names	CBFB; CBFbeta; PEBP2B; PEA2beta; PEBP2beta
Accession No.	Swiss-Prot:Q13951NCBI Gene ID:865
SDS-PAGE MW	22KD
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C


Application Details

Western blotting: 1:500 - 1:2000


Immunohistochemistry: 1:50 - 1:200

Immunofluorescence: 1:50 - 1:200

Images



Western blot analysis of extracts of various cell lines, using CBFB antibody.



Immunofluorescence analysis of U2OS cell using CBBF antibody. Blue: DAPI for nuclear staining.

Background

The protein encoded by this gene is the beta subunit of a heterodimeric core-binding transcription factor belonging to the PEBP2/CBF transcription factor family which master-regulates a host of genes specific to hematopoiesis (e.g., RUNX1) and osteogenesis (e.g., RUNX2). The beta subunit is a non-DNA binding regulatory subunit; it allosterically enhances DNA binding by alpha subunit as the complex binds to the core site of various enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers and GM-CSF promoters. Alternative splicing generates two mRNA variants, each encoding a distinct carboxyl terminus. In some cases, a pericentric inversion of chromosome 16 produces a chimeric transcript consisting of the N terminus of core-binding factor beta in a fusion with the C-terminal portion of the smooth muscle myosin heavy chain 11. This chromosomal rearrangement is associated with acute myeloid leukemia of the M4Eo subtype. Two transcript variants encoding different isoforms have been found for this gene.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.