

S-Tag Mouse Monoclonal Antibody

Catalog No: #T611

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	S-Tag Mouse Monoclonal Antibody
Host Species	Mouse
Clonality	Monoclonal
Isotype	IgG1
Purification	Affinity-chromatography
Applications	WB
Species Reactivity	Tag
Specificity	S- Tag antibody can recognize C-terminal, internal, and N-terminal S-tagged proteins.
Immunogen Type	Peptide-KLH
Immunogen Description	A synthetic peptide KETAAAKFERQHMS coupled to KLH.
Target Name	S-Tag
SDS-PAGE MW	N/A
Concentration	1.0mg/ml
Formulation	Mouse IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Application Details

WB: 1:3000~1:10000

Images



2ug S-Tag fusion protein+ Primary antibody dilution at
1γ • 1:5,000 2γ • 1:10,000

Background

S-tag is the name of an oligopeptide derived from pancreatic ribonuclease A (RNase A). If RNase A is digested with subtilisin, a single peptide bond is cleaved, but the resulting two products remain weakly bound to each other and the protein, called ribonuclease S, remains active although each of the two products alone shows no enzymatic activity. The N-terminus of the original RNase A, also called S-peptide, consists of 20 amino acid residues, of which only the first 15 are required for ribonuclease activity. This 15 amino acids long peptide is called S15 or S-tag. The amino acid sequence of the S-tag is: KETAAAKFERQHMS conjugated to KLH. S- Tag antibody can recognize C-terminal, internal, and N-terminal S-tagged proteins.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.