# Anti-DMPO Nitrone Adduct Catalog# SMC-189D

Size: 100ug

PO Box 55036 Cadboro Bay 3825 Cadboro Bay Rd, Victoria, BC V8N 4G0, Canada

# StressMarq Biosciences Inc.

Orders • sales@stressmarq.com

Tel: • +1 250 294 9065 Fax: • +1 250 294 9025

Email • info@stressmarq.com
Web • www.stressmarq.com

This product is for in vitro research use only and is not intended for use in humans or animals

Adduct Monoclonal Antibody  Clone  N1664A  Immunogen  5,5-dimethyl-2-(8-octanoic acid)-1-pyrrolone-N-oxide conjugated to Ovalbumin.  Host and Subclass  Mouse, IgG1  Cited Applications  Immuno-spin trapping, ELISA, WB, ICC/IF, IP (8)  Specificity  Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
CloneN1664AImmunogen5,5-dimethyl-2-(8-octanoic acid)-1-pyrrolone-N-oxide conjugated to Ovalbumin.Host and SubclassMouse, IgG1Cited ApplicationsImmuno-spin trapping, ELISA, WB, ICC/IF, IP (8)SpecificityDetects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.Species cross-reactivitySpecies IndependentFormatProtein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.Working dilution1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).Storage and-20°C; 1 year+; shipped on	Product	711.01.01.0
Immunogen  5,5-dimethyl-2-(8-octanoic acid)-1-pyrrolone-N-oxide conjugated to Ovalbumin.  Host and Subclass  Mouse, IgG <sub>1</sub> Cited Applications  Immuno-spin trapping, ELISA, WB, ICC/IF, IP (8)  Specificity  Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25μg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
acid)-1-pyrrolone-N-oxide conjugated to Ovalbumin.  Host and Subclass Mouse, IgG <sub>1</sub> Cited Applications Immuno-spin trapping, ELISA, WB, ICC/IF, IP (8)  Specificity Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution 1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25μg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on	Clone	
conjugated to Ovalbumin.  Host and Subclass Mouse, IgG1  Cited Applications Immuno-spin trapping, ELISA, WB, ICC/IF, IP (8)  Specificity Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution 1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25µg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on	Immunogen	
Host and Subclass       Mouse, IgG1         Cited Applications       Immuno-spin trapping, ELISA, WB, ICC/IF, IP (8)         Specificity       Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.         Species cross-reactivity       Species Independent         Format       Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.         Working dilution       1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).         Storage and       -20°C; 1 year+; shipped on		
Cited Applications Immuno-spin trapping, ELISA, WB, ICC/IF, IP (8)  Specificity Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution 1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on		conjugated to Ovalbumin.
WB, ICC/IF, IP (8)  Specificity  Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25μg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on	Host and Subclass	Mouse, IgG <sub>1</sub>
Specificity  Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross- reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on	Cited Applications	
DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on	C 'C' 'I	
DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25μg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on	Specificity	•
DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
cross-react with non-adducted proteins or DNA.  Species cross-reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25μg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
proteins or DNA.  Species cross- reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25μg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
Species cross- reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
reactivity  Format  Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10μg/mL (WB), 1-10μg/mL (ELISA), 10μg/mL (ICC), 25μg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		•
Format Protein G Purified. In TBS pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25µg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		Species Independent
pH7.4, with 0.05% sodium azide and 50% glycerol.  Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
azide and 50% glycerol.  Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25µg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on	Format	
Working dilution  1-10µg/mL (WB), 1-10µg/mL (ELISA), 10µg/mL (ICC), 25µg (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and  -20°C; 1 year+; shipped on		
(ELISA), 10µg/mL (ICC), 25ug (IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on		
(IP), see (8). For IP - DO NOT link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on	Working dilution	
link to matrix prior to use as this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on		(ELISA), 10μg/mL (ICC), 25ug
this causes large loss of reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on		
reactivity to DMPO adduct (see ref 8 for details).  Storage and -20°C; 1 year+; shipped on		link to matrix prior to use as
ref 8 for details).  Storage and -20°C; 1 year+; shipped on		
Storage and -20°C; 1 year+; shipped on		
stability cold packs or ambient	Storage and	-20°C; 1 year+; shipped on
, ,	stability	cold packs or ambient

#### Scientific Background

The formation of free radicals and other highly reactive oxygen species has been implicated in the pathogenesis of many disease states (1). The ability to identify these species is crucial, and spin trapping has accomplished

this goal. DMPO (5,5-dimethyl-1-pyrroline N-oxide) is one of the least toxic to cells and animals, and possesses convenient pharmacokinetics (uptake, distribution, metabolism and excretion) in biological systems (2-6).

Recent studies have determined that nitric oxide may substantially affect the quantitative determination of DMPO adducts, and therefore extra caution is required when studying generation of these species in the presence of nitric oxide or its radicals (1). DMPO adducts can be generated with protein and DNA radicals (7).

#### **Selected References**

- 1. Reszka K.J., et al. (2006) Nitric Oxide 15: 133-141.
- Ramirez D.C., Gomez-Mejiba S.E., and Mason R.P. (2007) Nat Protoc. 2(3): 512-522.
- Khan N., et al. (2003) Free Radic. Biol. Med 34:1473-1481.
- 4. Haseloff R.F., et al. (1997) FEBS Lett 418:73-75.
- 5. Schaefer C.F., Janzen E.G., West M.S., Poyer J.L., and Kosanke S.D. (1996) Free Radic. Biol. Med 21:427-436.
- 6. Anzai K., et al. (2003) Arch. Biochem. Biophys 415:251-256.
- 7. Free Radic Biol Med. 2009 April 1; 46(7): 853-865. doi:10.1016/j.freeradbiomed.2008.12.020.
- 8. Chatterjee S., Ehrenshaft, M., Bhattacharjee, S., Derterding, L.J., Bonini, M.G., Corbett, J., Kadiiska, M.B., Tomer K.B. and Mason, R.P. 2009 Free Radic. Med. and Biol. 46:454-461

### Certificate of Analysis

A 1/1000 dilution of SMC-189 was sufficient to detect the DMPO nitrone adducts of metmyoglobin when loaded at 100ng/lane by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody..

## Material Safety Data Sheet

### Anti-DMPO (Monoclonal Antibody) SMC-189

This product is for in vitro research use only and is not intended for use in humans or animals

The below information is believed to be correct but does not purport to be all inclusive and shall be used only as a guide. StressMarq shall not be held liable for any damage resulting from handling or from contact with the above product. See the Technical Specification, Packing Slip, Invoice, and Product Catalogue for additional terms and conditions of sale.

#### **Hazardous Ingredients**

The physical, chemical and toxicological properties of these components have not been fully investigated. It is recommended that all laboratory personnel follow standard laboratory safety procedures when handling this product. Safety procedures should include wearing OSHA approved safety glasses, gloves and protective clothing. Direct physical contact with this product should be avoided.

Known Hazardous Components
None

CAS Number
Percent

#### **Physical Data**

This product consists of mouse immunoglobulin in 50% glycerol and sodium azide shipped on gel packs. The physical properties of this product have not been investigated thoroughly.

#### Fire and Explosion Hazard and Reactivity Data

#### NOT APPLICABLE

#### **Toxicological Properties**

May be harmful by inhalation, ingestion, or skin absorption. The toxicological properties of this product have not been investigated thoroughly. Exercise due caution.

#### **Preventative Measures**

Wear chemical safety goggles and compatible chemical-resistant gloves. Avoid inhalation, contact with eyes, skin or clothing.

#### Spill and Leak Procedures

Observe all federal, state and local environmental regulations.

- Wear protective equipment.
- Absorb on sand or vermiculite and place in closed containers for disposal.
- Dispose or mix the material with a combustible solvent and burn in a chemical incinerator equipped with an afterburner and scrubber.

#### First Aid Measures

- If swallowed, wash out mouth with water, provided person is conscious. Call a physician.
- In case of skin contact, flush with copious amounts of water for at least 15 minutes. Remove contaminated clothing and shoes. If a rash or other irritation develops, call a physician.
- If inhaled, remove to fresh air. If breathing becomes difficult, call a physician.
- In case of eye contact, flush with copious amounts of water for at least 15 minutes while separating the eyelids with fingers. Call a physician.

Authorized: StressMarq Biosciences Inc. Creation Date: 10/01/2010