

## CD25 Antibody : B-B10 Azide free

HUMAN CD25 AZIDE FREE

**CATALOGUE N° :** 852.000.000

**CLONE :** B-B10

**ISOTYPE :** Mouse IgG1

**IMMUNISATION :** PHA activated T cells

**HYBRIDOMA :** Myeloma X63/AG.8653 x Balb/c spleen cells

**SPECIFICITY :** Recognises the Interleukin 2 Receptor a chain (IL-2 Ra), a 55 kDa protein

**WORKSHOP :** IV - V

**QUANTITY :** 200 µg / 200 µl

**PURIFICATION :** Ion exchange chromatography

**FORMAT :** Phosphate-buffered saline. Sterile-filtered through 0.22 µm and treated to remove endotoxins. Carrier and preservative free


**STORAGE :** Stable at +2-8°C for 12 months  
For longer storage, freeze aliquots

**APPLICATION :** Suitable for flow cytometry  
Inhibits Mixed Lymphocyte Culture  
Immunoprecipitation : not tested  
Western Blot : not tested  
Immunohistochemistry : not tested  
Recognises only membrane bound human IL-2 Ra

**OTHER :** Kd : 0.4 nM

**REFERENCES :** Engert, A. et al., *Int J Cancer*,1991; 49(3): 450-6.  
Groux, H. et al., *J Immunol.*,1998; 160(7): 3188-93.  
Herbelin, C. et al., *Bone Marrow Transplant*,1994; 13(5): 563-9.  
Herve, P. et al., *Blood*,1990; 75(4): 1017-23.  
Herve, P. et al., *Lancet*,1988; 2(8619): 1072-3.  
Heyll, A. et al., *Dtsch Med Wochenschr.*,1991; 116(22): 852-6.

Mousson, C. et al., *Transplant Proc.*, 1995; 27(2): 1727-8.  
Nakatani, T. et al., *Protein Eng.* 1994; 7(3): 435-43.  
Nevill, T. J. et al., *Blood*, 1998; 92(6): 1910-7.  
Oppmann, B. et al., *Immunity*, 2000; 13(5): 715-25.  
Ozsahin, H. et al., *Blood*, 1998; 92(8): 2719-24.  
Racadot, E. et al., *Bone Marrow Transplant*, 1995; 15(5): 669-77.  
Racadot, E. et al., *Transplant Proc.*, 1995; 27(2): 1672-3.  
Regeer, R. R. and D. Markovich, *Am J Physiol Cell Physiol.*, 2004; 287(2): C365-72.  
Tazzari, P. L. et al., *Transplantation*, 1992; 54(2): 351-6.  
Tiberghien, P. et al., *Transplantation*, 1991; 52(3): 475-80.  
Wijdenes, J. et al., *Dev Biol Stand.*, 1990; 71: 103-11.  
Zambello, R. et al., *Blood*, 1997; 89(1): 201-11.



*A typical staining pattern with the B-B10 monoclonal antibody of PHA activated lymphocytes*