

### Anti-SARS-CoV-2 Nucleocapsid Protein (NCP), Mouse-Mono (Clone # 6)

Cat No. # SNCP1F

Package : 25 µg

<b>Host</b>	Mouse
<b>Immunogen</b>	Recombinant SARS-CoV-2 Nucleocapsid protein (NCP)
<b>Clonality</b>	Monoclonal
<b>Purification</b>	Protein G purified
<b>Application</b>	WB, ELISA
<b>Reactivity</b>	SARS Coronavirus 2

#### PROPERTIES

<b>Form</b>	Liquid
<b>Buffer</b>	1×PBS (pH 7), 0.09% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	0.5 mg/mL (Please refer to the vial label for the specific concentration)
<b>Conjugation</b>	Unconjugated
<b>Note</b>	For Research use. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


#### APPLICATION

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	Recombinant SARS-CoV-2 Nucleocapsid Protein
<b>ELISA</b>	1 µg/mL	Recombinant SARS-CoV-2 Nucleocapsid Protein

\*Application Note: Optimal dilutions/concentrations should be determined by the researcher.

### DATA IMAGES


#### Western Blot



#### Detection of recombinant SARS-CoV-2 NCP by Western Blot.

Western blot shows recombinant SARS-CoV-2 NCP. PVDF membrane was detected with 1 µg/mL of SARS-CoV-2 nucleocapsid protein (NCP) monoclonal antibody (Catalog # SNCP1F) followed by Anti-IgG (H+L chain) (Mouse) pAb-HRP (Code No. 330). A specific band was detected for NCP at approximately 55 kDa (as indicated). This experiment was conducted under reducing conditions.

#### Indirect ELISA



#### Detection of recombinant SARS-CoV-2 NCP by indirect ELISA.

Indirect ELISA detection of diluted at 125-4000 ng/mL recombinant SARS-CoV-2 nucleocapsid protein using SARS-CoV-2 nucleocapsid protein (NCP) monoclonal antibody (Catalog # SNCP1F) as capture antibody at concentration of 1 µg/mL, and Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed (ab150117) was diluted at 1:5000 and used to detect the primary antibody.