

INSTRUCTION MANUAL

Quick-gDNA™ Blood MicroPrep

Catalog Nos. **D3070** & **D3071**

Highlights

- Quick purification of high quality DNA from *whole blood*, *plasma*, and *serum* in less than 15 minutes using innovative *Fast-Spin* column technology.
- Compatible with commonly used anticoagulants (i.e., EDTA, heparin, citrate).
- Unique extraction technology <u>excludes</u> the use of Proteinase K and organic denaturants.
- Isolated DNA is ideal for PCR, endonuclease digestion, bisulfite conversion/methylation detection, sequencing, genotyping, etc.

Contents

Product Contents	1
Specifications	1
Product Description	2
Buffer Preparation	3
Protocol	3
Troubleshooting	4
Ordering Information	4
Related Products	5

For Research Use Only Ver. 1.0.1

Satisfaction of all Zymo Research products is guaranteed. If you should be dissatisfied with this product please call 1-888-882-9682.

Use the *Quick-gDNA*™ Blood MiniPrep (D3072, D3073) for

For high-throughput purification (96-well, 5 µg DNA/well) use

Blood (D3075, D3076, D3077).

the ZR-96 Quick-gDNA™

the recovery of up to 25 µg/prep DNA from small

samples.

Product Contents

Quick-gDNA™ Blood MicroPrep (Kit Size)	D3070 (50 preps.)	D3071 (200 preps.)	Storage Temperature
Genomic Lysis Buffer*	50 ml	2x 100 ml	Room Temp.
DNA Pre-Wash Buffer**	15 ml	50 ml	Room Temp.
g-DNA Wash Buffer	50 ml	100 ml	Room Temp.
DNA Elution Buffer	10 ml	2x 10 ml	Room Temp.
Zymo-Spin™ IC Columns	50	200	Room Temp.
Collection Tubes	100	400	Room Temp.
Instruction Manual	1	1	-

Note - Integrity of kit components is guaranteed for up to one year from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide the highest performance and reliability.

Specifications

- Sample Sources Whole blood, plasma, or serum from humans, mice, rats, etc.
- **DNA Purity** High-quality DNA is eluted with **DNA Elution Buffer** or water. DNA is especially well suited for PCR and other downstream applications. A_{260}/A_{280} >1.8
- Size Limits Capable of recovering genomic DNA up to and above 40 kb. In most instances, mitochondrial DNA and viral DNA (if present) will also be recovered.
- DNA Recovery Up to 5 µg total DNA is eluted into ≥10 µl (6 µl minimum) DNA
 Elution Buffer or water. Human whole blood will typically
 yield 3-7 µg DNA per 100 µl blood sampled.
- Product Detergent Tolerance ≤5% Triton X-100, ≤5% Tween-20, ≤5% Sarkosyl,
 ≤0.1% SDS.
- Equipment Needed Microcentrifuge

Note - TM Trademarks of Zymo Research Corporation. This product is for research use only and should only be used by trained professionals. It is not intended for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

^{*} $\underline{\textit{Recommended}}$: Add beta-mercaptoethanol to 0.5%(v/v) i.e., 250 μ l per 50 ml or 500 μ l per 100 ml.

^{**} A precipitate may have formed in the DNA Pre-Wash Buffer during shipping. To completely resuspend the buffer, incubate the bottle at 30-37 °C for 30 minutes and mix by inversion. DO NOT MICROWAVE.

Product Description

The **Quick-gDNA™ Blood MicroPrep** is a simple procedure for the rapid isolation of total DNA (e.g., genomic, mitochondrial, viral) from a variety of biological sample sources. This product has been optimized for maximal recovery of ultra-pure DNA without RNA contamination and is compatible with whole blood (fresh or stored), serum, and plasma.

For processing, simply add the specially formulated **Genomic Lysis Buffer** to a sample, vortex, and transfer the mixture to the supplied **Zymo-Spin™ Column**. There is no need for organic denaturants or Proteinase K digestion because of the unique chemistries featured in the kit. Instead, the product features *Fast-Spin* technology to yield high-quality, purified DNA in just minutes (see below). PCR inhibitors are effectively removed during the purification process. DNA purified using the *Quick-gDNA™* **Blood MicroPrep** is suitable for PCR, nucleotide blotting, DNA sequencing, restriction endonuclease digestion, bisulfite conversion/methylation analysis, and other downstream applications.

Sample

+
Genomic Lysis Buffer

Spin
Wash
Elute

Ultra-pure DNA is ideal for...

- ✓ PCR
- ✓ Restriction Endonuclease Digestion
- ✓ Genotyping
- ✓ Bisulfite Conversion & Methylation Analysis

Zymo Supplier Supplier S

High yield/quality DNA is successfully isolated from porcine whole blood using the *Quick-gDNA™* Blood MiniPrep (D3072). Equivalent amounts (100 µl) of blood were processed without Proteinase K using the *Quick-gDNA™* Blood MiniPrep in half the time as compared to the kits from suppliers Q and S. Equal volumes of eluted DNA were then analyzed (in duplicate) in a 0.8% (w/v) TAE/agarose/ethidium bromide gel. The size marker "M" is a 1 kb ladder (Zymo Research).

The ZymoBead™ Genomic DNA Kit (D3004, D3005) and ZR Serum DNA Kit™ (D3013) are recommended for scaleable DNA yields >25 µg/prep. Both feature silica beads instead of a spin column.

Zymo Research offers the following for rapid, precise DNA methylation detection...

- 1.) **EZ DNA Methylation™ Kit** (D5001, D5002, D5003, D5004)
- 2.) **EZ DNA Methylation- Gold™ Kit** (D5005, D5006, D5007, D5008)
- 3.) **EZ DNA Methylation- Direct™ Kit** (D5020, D5021, D5022, D5023)

For **Technical Assistance**, please contact 1-888-882-9682 or E-mail tech@zymoresearch.com.

Buffer Preparation

Recommended: Add beta-mercaptoethanol (user supplied) to the **Genomic Lysis Buffer** to a final dilution of 0.5%(v/v) i.e., 250 μl per 50 ml *or* 500 μl per 100 ml.

Protocol

The following is for the purification of DNA from 50 µl whole blood, serum or plasma (the volumes can be adjusted <u>up to</u> 100 µl (max.) depending on your requirements). Fresh, frozen, or preserved blood (in EDTA, citrate, or heparin) can be used. If material cannot be processed immediately, the sample can be "stabilized" for later processing (as noted below) although the immediate processing of blood samples is recommended.

 Add 200 µl of Genomic Lysis Buffer to 50 µl of blood, serum, or plasma (4:1). Mix completely by vortexing 4-6 seconds, then let stand 5-10 minutes at room temperature.¹

Note: Add 200 μl Genomic Lysis Buffer to all samples <50 μl. For samples larger than 50 μl, add a proportional amount (4:1) of Genomic Lysis Buffer (e.g., Add 800 μl Genomic Lysis Buffer to 200 μl blood).

- 2. Transfer the mixture to a **Zymo-SpinTM IC Column²** in a **Collection Tube**. Centrifuge at $10,000 \times g$ for one minute. Discard the **Collection Tube** with the flow through.
- 3. Transfer the **Zymo-Spin™ Column** to a new **Collection Tube**. Add 200 µl of **DNA Pre-Wash Buffer** to the spin column. Centrifuge at 10,000 *x g* for one minute.
- 4. Add 500 μ l of **g-DNA Wash Buffer** to the spin column. Centrifuge at 10,000 x q for one minute.
- 5. Transfer the spin column to a clean microcentrifuge tube. Add ≥10 µl **DNA Elution Buffer** or water to the spin column³. Incubate 2-5 minutes at room temperature and then centrifuge at top speed for 30 seconds to elute the DNA. The eluted DNA can be used immediately for molecular based applications or stored ≤-20°C for future use.

Delayed Processing (Stabilization) of Blood Samples: The immediate processing of blood with this kit is recommended. However, if blood cannot be processed immediately, samples can be "stabilized" in Genomic Lysis Buffer for processing at a later time. To do this, add *four* volumes of Genomic Lysis Buffer to *each* volume of whole blood (4:1), then vortex. Blood samples mixed with Genomic Lysis Buffer can be stored at room temperature for 1-2 weeks, 0-4°C for 1-2 months, -20°C for 6 months to a year, or <-70°C for many years. Samples stored at ≤4°C should reach room temperature prior to processing. Begin at Step 2 in the standard protocol (above) when purifying DNA from blood samples stabilized in Genomic Lysis Buffer.

Notes:

- ¹ For the inclusion of <u>small DNAs</u> from serum, add 0.3 volumes isopropanol to the mixture. (For example, to a 800 µl mixture of serum and Genomic Lysis Buffer add 240 µl isopropanol.)
- ² The column capacity is 800 ul.
- ³ Elution of DNA from the column is dependent on pH and temperature. If water is used, ensure the pH is >6.0. Also, the total yield may be improved by eluting the DNA with Elution Buffer or water preequilibrated to 60-70°C.

Troubleshooting:

- 1. <u>DNA degradation</u>: Check for DNase contamination. All reagents supplied with the *Quick-gDNA*TM **Blood MicroPrep** are DNase-free. However, DNase contamination could result during the processing of some samples. Check pipets, pipet tips, microcentrifuge tubes, etc., and exercise the appropriate precautions during the DNA purification procedure.
- 2. <u>DNA is not performing well in subsequent experiments</u>: Ensure the correct volume of **Genomic Lysis Buffer** has been added to the sample. Also, make sure all centrifugation steps are completed for the indicated times and speeds (rcfs). Failure to do so may result in incomplete washing, which may cause salts to be eluted with the DNA affecting quantitation and subsequent experiments including enzymatic processes like PCR.
- 3. RNA contamination: The buffers in this kit are designed to efficiently hydrolyze and remove RNA during the DNA purification procedure.
- 4. <u>Avian Blood:</u> Reduce input to 2-5 μl when working with avian blood. Add 200 μl of **Genomic Lysis Buffer** to the sample. Thoroughly mix the sample by vortexing intermittently for 5 minutes. Transfer to the **Zymo-Spin™ Column** and proceed with step 2 of the protocol.

Ordering Information

Product Description	Cat. No.	Kit Size
<i>Quick-gDNA</i> ™ Blood MicroPrep	D3070 D3071	50 preps. 200 preps.
<i>Quick-gDNA</i> ™ Blood MiniPrep	D3072 D3073	50 preps. 200 preps.
<i>Quick-gDNA</i> ™ Blood MidiPrep	D3074	25 preps.
ZR-96 <i>Quick-gDNA</i> ™ Blood	D3075 D3076 D3077	2x 96 well 4x 96 well 10x 96 well

For Individual Sale	Cat. No.	Amount
Genomic Lysis Buffer	D3004-1-50 D3004-1-100	50 ml 100 ml
DNA Pre-Wash Buffer	D3004-5-15 D3004-5-30 D3004-5-50	15 ml 30 ml 50 ml
g-DNA Wash Buffer	D3004-2-50 D3004-2-100	50 ml 100 ml
DNA Elution Buffer	D3004-4-10	10 ml
Zymo-Spin™ IIC Columns	C1011-50 C1011-250	50 250
Collection Tubes	C1001-50 C1001-500 C1001-1000	50 500 1,000

THE Epigenetics COMPANYTM

Popular Products From Zymo Research

Product	Description	Kit Size	Catalog No.
Troduct	·	(Preps)	(Format)
	Fragment DNA Purification		
DNA Clean & Concentrator™-5	Clean and concentrate up to 5µg DNA into ≥6 µl elution volume in as little as 2 minutes with no wash residue carryover.	50 200 50 200	D4003 (uncapped) D4004 (uncapped) D4013 (capped) D4014 (capped)
DNA Clean & Concentrator™-25	Clean & concentrate 25 µg of DNA into ≥25 µl elution volume in as little as 2 minutes with no wash residue carryover.	50 200 50 200	D4005 (uncapped) D4006 (uncapped) D4033 (capped) D4034 (capped)
ZR-96 DNA Clean & Concentrator™-5	Quick (15 minute), high-output recovery of up to 5 µg pure DNA into 10-15 µl minimum elution volume allows for highly concentrated DNA.	2x96 4x96	D4023 D4024
Genomic DNA Clean & Concentrator™	Quick (5 minute) clean-up of up to 10 µg high molecular weight DNA (≤200 kb) from any enzymatic reaction or impure preparation without precipitations.	25 100	D4010 (capped) D4011 (capped)
Zymoclean™ Gel DNA Recovery Kit	Purify DNA from high and low-melting agarose gels in minutes	50 200 50 200	D4001 (uncapped) D4002 (uncapped) D4007 (capped) D4008 (capped)
ZR-96 Zymoclean™ Gel DNA Recovery Kit	High-throughput DNA purification from high and low-melting agarose gels.	2x96 4x96	D4021 D4022
Zymoclean™ Large Fragment DNA Recovery Kit	Purify high molecular weight DNA (≤200 kb) from high and low-melting agarose gels in minutes	25 100	D4045 (capped) D4046 (capped)
OneStep™ PCR Inhibitor Removal Kit	Fast, one step procedure for removal of PCR inhibitors such as polyphenolics, humic/fulvic acids, melanin, etc. for successful PCR and other downstream applications.	50 2x96	D6030 D6035
	Plasmid DNA Purification		
Zyppy™ Plasmid Miniprep Kit	Pellet-Free™ plasmid DNA purification in less than 10 minutes. Recover up to 25 µg DNA in as low as 30 µl.	50 100 400	D4036 D4019 D4020
Zyppy™-96 Plasmid Miniprep	The fastest and simplest high-throughput method for plasmid purification.	2x96 4x96 8x96	D4041 D4042 D4043
Zyppy™ Plasmid Midiprep Kit	Pellet-Free™ plasmid DNA purification in 15 minutes in a 150 µl minimum elution volume).	25 50	D4025 D4026
ZR Plasmid MiniPrep™ Classic	Plasmid DNA purification in minutes: (alkaline lysis/spin column format for low 30 µl elution volume).	50 100 400	D4036 D4019 D4020
	Genomic DNA Purification		
<i>Quick-gDNA</i> ™ MiniPrep	Easy purification of genomic DNA from whole blood, plasma, serum, body fluids, buffy coat, lymphocytes, tissue, swabs or cultured cells in as little as 15 minutes without the use of Proteinase K or organic denaturants.	50 200 50 200	D3006 (uncapped) D3007 (uncapped) D3024 (capped) D3025 (capped)
ZR-96 <i>Quick-gDNA</i> ™	Simple, high throughput (96-well) purification of DNA from whole blood, plasma, serum, body fluids, buffy coat, lymphocytes, tissue, swabs, or cultured cells in about 30 minutes.	2x96 4x96 10x96.	D3010 D3011 D3012
ZR-Genomic DNA™- Tissue MiniPrep	For high quality DNA purification from solid tissues (e.g., tail snips, ear punches, adipose tissue, etc.), body fluids, cultured cells, buccal cells, FFPE tissues, hair, and other biological sources using Proteinase K and Fast.	50 200	D3050 D3051
Environmental DNA Purification Kits	Unique BashingBead™ technology allows isolation of DNA from samples refractory to conventional lysis procedures including tough-to-lyse tissues, soil samples, feces, plants, seeds, insects, bacteria, yeast, filamentous fungi, unicellular and filamentous algae, and protozoa		Visit website for a comprehensive list

Please visit our website to see our complete line-up of products.