

ZymoScript™ One-Step RT-qPCR Kit

Simple and sensitive qPCR detection of RNA targets

Cat. No. R3014 (100 Reactions)

Storage: -20 °C

Product Information

Features:

- Versatile and fast RT-qPCR compatible with both SYBR Green and TaqMan-based assays
- Efficient cDNA synthesis coupled with robust amplification
- Reduced non-specific product formation
- Ideal for pathogen detection and gene expression analysis
- Tracking dye to monitor pipetting errors

Description:

The **ZymoScript™ One-Step RT-qPCR Kit** contains all the necessary components for fast reverse transcription and robust qPCR amplification in a single reaction. Simply add RNA template and primers to achieve streamlined RT-qPCR in a single tube.

ZymoScript™ One-Step RT-qPCR Kit is compatible with both SYBR Green and TaqMan probe-based assays for ultimate flexibility. The provided Fluor Dye solution can be added to the RT-qPCR reaction for SYBR Green-based detection. Alternatively, TaqMan probes can be added for multiplex RT-qPCR analysis. cDNA synthesis and robust target amplification can be achieved in 1.5 hour. The provided enzyme mix contains optimized elements for maximum specificity during each reaction step.

The resulting RT-qPCR product contains an "A" overhang to allow for TA-cloning.

Product Contents:

	Catalog Number	R3014 (100 reactions)	Storage Temp.
Reaction Buffer (4X)	R3014-1-500	1 x 500 µl	-20 °C
Enzyme Mix (20X)	R3014-2-100	1 x 100 µl	-20 °C
Fluor Dye (20X)	R3014-3-100	1 x 100 µl	-20 °C
MgCl ₂ (25 mM)	R3014-4-500	1 x 500 µl	-20 °C
DNase/RNase-Free Water	W1001-1	1 x 1 ml	RT

Note: Fluor Dye must be protected from light.

Storage:

Store at -20 °C. Minimize exposure to light.

Suggested Reaction Setup (20 µl):

Reagent	Volume	Final conc.
Reaction Buffer (4X)	5.0 µl	1X
Enzyme Mix (20X)	1.0 µl	1X
Forward Primer (10 µM)	0.8 µl	0.4 µM
Reverse Primer (10 µM)	0.8 µl	0.4 µM
Fluor Dye (20X)*	1.0 µl	1X
*For SYBR Green assays only		
Template RNA	variable	0.1 pg to 5 µg total
DNase/RNase-Free Water	to 20 µl	–
Total volume	20 µl	

Notes:

- Only add Fluor Dye if performing SYBR Green-based fluorescent detection.
- MgCl₂ is present in the reaction buffer. Its concentration in the final RT-qPCR reaction is 1.6 mM, but can be increased up to 4 mM using the provided 25 mM MgCl₂.

Suggested Conditions For RT-qPCR:

1) Primer annealing & cDNA synthesis ¹	55 °C	10 min.
2) Initial denaturation	95 °C	10 min.
3) Denaturation	95 °C	20 sec.
4) Annealing/Extension ²	60 °C	60 sec. for ≤ 1kb
5) Plate Read	30-45 cycles	
6) Optional: Melt Curve	60-95 °C	
*For SYBR Green assays only		

¹ For the cDNA synthesis of long transcripts, extend this step up to 45 min.

² Optimal annealing temperature may vary based on primer design. Adjust the temperature if necessary. Add 60 seconds to the extension time for each additional kb over 1 kb.

- If cloning after RT-qPCR is desired, include a final extension step of 72°C for 5 minutes.

Related Products:


Product	Cat. Number	Description
Quick-RNA™ Kits	R1050	RNA isolation from wide range of cell and tissue samples
Direct-zol™ RNA Kits	R2050	RNA extraction from samples in TRI Reagent® or similar
DNase I Set	E1010	DNase I enzyme and digestion buffer
ZymoScript™ RT PreMix Kit	R3012	RT kit, ready to use PreMix formulation. Use for 2-step RT-qPCR

Continued on reverse side...

Additional Information

ZymoScript™ One-Step RT-qPCR Analysis:

The ZymoScript™ One-Step RT-qPCR Kit enables highly sensitive target detection compared to other suppliers



RNA was isolated from HeLa cells. Values are represented as mean (+/- SE) cycle threshold (C_t) of 3 RT-qPCR technical replicates. Targets 1, 2 and 3 are GAPDH, ACTB, and TBP, respectively.

FAQ:

1. What is the minimum/maximum amount of RNA template I should use?

The amount of total RNA required may vary depending on the expression level of the target transcripts. In general, we recommend using 0.1 pg - 5 µg of input RNA.

2. Will the blue tracking dye interfere with qPCR fluorophore channels and affect the signal of probe/fluorescent dyes?

The dye used in the ZymoScript™ One-Step RT-qPCR Kit has been thoroughly tested in downstream analysis. At the provided concentration, it will not affect fluorescent signal readings during qPCR.

3. How is the performance of the ZymoScript™ One-Step RT-qPCR Kit with low-quality RNA samples?

Lower signal can be expected with degraded and impure RNA samples. We recommend using *Quick-RNA*™ or *Direct-zol*™ Kits (see **Related Products**) for high quality RNA extraction.

4. Can I use oligo dT or random hexamer primers?

No, only use gene-specific primers when using ZymoScript™ One-Step RT-qPCR Kit.

5. Can I assemble the RT-PCR reaction at room temperature?

Yes, the reaction can be assembled at room temperature. The enzymes are only activated at higher temperatures, ensuring maximum specificity.

6. Can I store the ZymoScript™ One-Step RT-qPCR Kit at Room Temperature or 4 °C?

We do not recommend storing the kit or kit components at room temperature or at 4 °C for prolonged periods of time.

7. How can I ensure that there is no genomic DNA contamination in my RT reaction?

Samples can be treated with DNase I to eliminate DNA contamination. DNase I is included in the *Quick-RNA*™ or *Direct-zol*™ Kits recommended for RNA extraction. Alternatively, the DNase I set (see **Related Products**) can be purchased separately.

8. Do I need to add RNase inhibitors?

No, RNase inhibitors are already included in the enzyme mix to prevent unwanted RNA degradation during reverse transcription.

9. Should I use SYBR Green or a TaqMan probe-based assay?

For single target detection, SYBR Green-based assays offer a more affordable solution than TaqMan probe-based assays. For SYBR Green-based assays, add 1 µl of Fluor Dye solution (20X) to each 20 µl RT-qPCR reaction. For TaqMan probe-based assays we recommend adding TaqMan probes at a final concentration of 0.2 µM. Do not add Fluor Dye solution to the RT-qPCR reaction when performing TaqMan probe-based assays.

10. Does ZymoScript™ One-Step RT-qPCR Kit contain a passive reference dye?

No, a passive reference dye is not included. This allows to maximize the number of targets for multiplex assays.

11. What is the maximal amplicon length that can be achieved using the ZymoScript™ One-Step RT-qPCR Kit?

Although this product performs the best with relatively short amplicons (50 bp – 1 Kb), it has been successfully used to amplify regions up to 5 Kb long.

Trademarks and Disclaimers:

This product is for research use only and should only be used by trained professionals. It is not intended for use in diagnostic procedures. Some reagents included with this kit are irritants. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

The Polymerase Chain Reaction (PCR) process is covered by U.S. Patent: #4,683,195;4,683,202 assigned to Hoffmann-La Roche. Patents pending in other countries. No license under these patents to use the PCR process is conveyed expressly or by implication to the purchaser by the purchase of Zymo Research's products. Further information on purchasing licenses to practice the PCR process can be obtained from the director of Licensing at Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California 94404 or at Roche Molecular Systems, Inc., 1145 Atlantic Avenue, Alameda, California 94501.

SYBRgreen® is a registered trademark of Life Technologies Corporation. ZymoScript™ is a trademark of Zymo Research Corporation.