



Anti-human VLDL receptor Antibody Monoclonal, VR-VP1-5C3

ORDERING INFORMATION

Catalog Number: BML030

Lot Number:

Size: 50 µg

Formulation: 0.2 µm filtered PBS solution

Storage: -80°C

Specificity: human VLDL receptor

Immunogen: synthetic peptide

Ig Type: IgG2b

Application: Western blot

Flowcytometry

Immunohistochemistry

Preparation

Produced in mice immunized with synthetic peptides, amino acid residue 1-18 (GRKAKCEPSQFQCTNGRC), which is corresponding to the NH₃-terminus of human very low-density lipoprotein (VLDL) receptor (VLDL-R). VLDL-R specific IgG was purified from mouse ascites fluid with a protein A-Sepharose.

Formulation

0.2 µm filtered PBS solution

Storage

IgG in PBS solution are stable for twelve months from the date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles.

Specificity

This antibody has been selected for its ability to bind for human VLDL-R expressed in CHO cells (*Idl-A7*). No cross-reactivity with human LDL receptor and apoER2 receptor was confirmed (see ref. 1).

Additional Applications

Western Blot - This antibody can be used at 1.0 µg/mL for western blot analysis (1).

Flow cytometry - This antibody can be used as a 1st antibody for immunohistochemistry. Please see the references (2) and (3) for details.

Immunohistochemistry - This antibody can be used for immunohistochemistry (4).

Optimal dilutions should be determined by each laboratory for each application.

References

- (1) Kosaka et al., Evidence of macrophage form cell formation by very low-density lipoprotein receptor: Interferon-γ inhibition of very low-density lipoprotein receptor expression and form cell formation in macrophages. *Circulation*, 2001;103:1142-1147.
- (2) Iwasaki et al., The important role for BVLDLs binding at the forth cysteine of first ligand-binding domain in the low-density lipoprotein receptor. *J Hum Genet*, 2004;49:622-628.
- (3) Iwasaki et al., Deficiency of the very low-density lipoprotein (VLDL) receptors in streptozotocin-induced diabetic rats: insulin dependency of the VLDL receptor. *Endocrinology*, 2005;146:3286-3294.
- (4) Motoi et al., Apolipoprotein E receptor 2 is involved in neurotic plaque formation in APP sw mice. *Neurosci Lett*, 2004;368:144-147.

FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

BML, Inc.

1361-1 Matoba, Kawagoe, Saitama 350-1101, Japan
TEL +81-49-232-0440/FAX +81-49-232-5480