

Anti-Laminin γ2 N-terminal fragment,


Human, Mouse-Mono (Clone P2H)

Catalog NO. FDV-0025

Research use only, not for human or animal therapeutic or diagnostic use. This product has been commercialized under the license from Yokohama City University.

Product Background

Laminins, which consist of three subunits called α , β and γ chains, are major cell adhesive components of extracellular matrix, especially basement membrane (BM). The laminin family is constituted of over 15 isoforms, expressed in a tissue-specific manner and plays differential roles in each tissue. Laminin γ 2 chain is a subunit of laminin-332 isoform, formerly laminin-5, but it is frequently overexpressed as a monomer form or the β 3- γ 2 heterodimer in invasive cancers. Laminin γ 2 is a protein of approximately 150 kDa and cleaved at its short arm mainly by bone morphogenic protein-1 (BMP1) in cancer cells, releasing an N-terminal proteolytic fragment (γ 2pf) of 45 kDa. The short arm and γ 2pf are also cleaved by MMPs and serine proteinases, releasing further small N-terminal fragments. Recently, Dr. Kaoru Miyazaki and co-workers generated a novel mouse monoclonal antibody against γ 2pf, clone P2H, which can efficiently detect invasive cancer cells. Clone P2H recognizes laminin γ 2 chain of laminin-332 on normal epithelial basement membranes, while in malignant cancer tissues it strongly immunostains the cytoplasm of cancer cells at invasion fronts. Thus, P2H is an important antibody capable of highly detecting the tumor invasion marker laminin γ 2 chain in cancer tissues. It is useful for pathological assessment of invasiveness and malignancy in various types of human cancers. P2H is applicable to not only immunohistochemistry, but also immunoblotting, ELISA and immunoprecipitation.





Description

Catalog Number : FDV-0025 Format : Mouse ascites Volume : 100 µL Formulation : Ascites without any additives Host Species and Clonality : Mouse monoclonal Isotype and Subclass : IgG2a Purification : No purification Lot Number : see vial label Specificity : Human, mouse, other species not tested yet Antigen : recombinant human γ 2pf containing domain IV + V Epitope : γ 2 domain V, NE2 Storage : For short-term storage, -20°C. For long-term storage, -80°C storage is preferable. Avoid repeated freeze-thaw cycles and avoid storage at 4°C.

Application

- Western blotting under both reducing, and non-reducing conditions
- Immunohistochemistry
- ELISA

Recommended usage

- Western blotting
- ELISA


1/1,000-1/10,000

- Immunohistochemistry

Optimal dilutions should be determined by the researcher.

Optimal conditions should be determined by the researcher.

Application examples



30

24

P2H

VMRC-LCP Medium

150

105 🗕

45 -

30 -

22 -

15

P2H

Immunohistochemistry : Normal mammary gland and breast cancer

Upper panel : Normal mammary gland Lower panel : Breast cancer

P2H stains basement membranes surrounding mammary glands in normal tissue, whereas in a breast cancer tissue cytoplasmic accumulation of γ 2pf or its fragments is detected with P2H.

Immunohistochemistry : Normal skin and skin cancer

Upper panel : Normal skin tissue Lower panel : Skin cancer tissue

Although P2H stains epithelial basement membranes in normal skin, in a skin cancer tissue it strongly stains invading cancer cells or their cytoplasm.

Immunohistochemistry : Lung cancers - SCC


Right upper panel : Stained by P2H (anti-γ2pf) Right lower panel : Stained by BG5 (anti-α3A, cat. No. FDV-0024)


Although BG5 detected basement membrane (BM)-like structure surrounding tumor cells, P2H detected not only BMs but also the cytoplasm of cancer cells.

$Immunohistochemistry: Lung\ SCC\ and\ ADC$


Upper two panels : Invasive lung SCC Lower two panels : Invasive lung ADC


P2H strongly detected cytoplasm of cancer cells at invasion fronts.











Reference

1. Miyazaki et al., Cancer Sci., 107, 1909-1918 (2016) Highly sensitive detection of invasive lung cancer cells by novel antibody against amino-terminal domain of laminin $\gamma 2$ chain.

luccu prou			
Catalog No.	Product name	Target	Application
FDV-0023	Anti-Laminin α3B, Human, Mouse-Mono (F7)	Laminin a3B	IHC, WB, IP, ELISA
FDV-0024	Anti-Laminin α3A, Human, Mouse-Mono (BG5)	Laminin a3A	IHC, WB, IP, ELISA
FDV-0025	Anti-Laminin γ2 N-terminal fragment,	Laminin y2	IHC, WB, ELISA
	Human, Mouse-Mono (P2H)	N-terminal fragments	
FDV-0026	Anti-Laminin 511, Human, Mouse-Mono (12D)	Trimeric Lm511 structure	IHC, WB, IP, ELISA

Related products

Disclaimer/免責事項

This product has been commercialized by Funakoshi Co., Ltd. based on the results of academic research, and the advertisement text, figures and manuals (hereinafter "Product information") have been prepared based on published research reports on December, 2018. The academic interpretation at the time of creation of the Product Information may change in accordance with future developments in the relevant research field and expansion of various scientific findings, and the latest version and certainty of the Product Information are not guaranteed. The specifications of this product and the Product Information are subject to change without notice. Please contact us for the latest information.

本製品は学術研究成果を基にフナコシ株式会社が製品化したもので、 2018年12月時点における公開研究報告を基に広告文章およびマニュ アル(以下、製品資料)を作成しています。今後の当該研究分野の発 展および各種学術知見の拡大にともない、製品資料作成時の学術的 解釈が変更になる可能性があり、最新性・確実性を保証するものでは ありません。また、本製品の仕様および製品資料を予告なく変更する 場合がございます。最新の情報に関しましては、弊社までご確認いた だきますようお願い申し上げます。





[ver. 2023/07]

Download the latest datasheet from www.funakoshi.co.jp www.funakoshi.co.jp/exports

(Japanese) (English)